


This article was downloaded by: [Uppsala universitetsbibliotek]

On: 30 December 2013, At: 23:57

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Synthetic Communications: An International Journal for Rapid Communication of Synthetic Organic Chemistry

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/lcyc20>

Green and Catalyst-Free One-Pot Synthesis of Anthranilamide Schiff Bases: An Approach Toward Sirtinol

Seyed Mostafa Ebrahimi ^a, Mohammad Mahdavi ^a, Saeed Emami ^b, Mina Saeedi ^a, Mehdi Asadi ^a, Loghman Firoozpour ^c, Mehdi Khoobi ^a, Kouros Divsalar ^d, Abbas Shafiee ^a & Alireza Foroumadi ^{a c}

^a Department of Medicinal Chemistry, Faculty of Pharmacy and Pharmaceutical Sciences Research Center, Tehran University of Medicinal Sciences, Tehran, Iran

^b Department of Medicinal Chemistry and Pharmaceutical Sciences Research Center, Faculty of Pharmacy, Mazandaran University of Medical Sciences, Sari, Iran

^c Drug Design and Development Research Center, Tehran University of Medicinal Sciences, Tehran, Iran

^d Neuroscience Research Center, Institute of Neuropharmacology, Kerman University of Medicinal Sciences, Kerman, Iran

Accepted author version posted online: 28 Oct 2013. Published online: 27 Dec 2013.

To cite this article: Seyed Mostafa Ebrahimi, Mohammad Mahdavi, Saeed Emami, Mina Saeedi, Mehdi Asadi, Loghman Firoozpour, Mehdi Khoobi, Kouros Divsalar, Abbas Shafiee & Alireza Foroumadi (2014) Green and Catalyst-Free One-Pot Synthesis of Anthranilamide Schiff Bases: An Approach Toward Sirtinol, *Synthetic Communications: An International Journal for Rapid Communication of Synthetic Organic Chemistry*, 44:5, 665-673, DOI: [10.1080/00397911.2013.833627](https://doi.org/10.1080/00397911.2013.833627)

To link to this article: <http://dx.doi.org/10.1080/00397911.2013.833627>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions

and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

GREEN AND CATALYST-FREE ONE-POT SYNTHESIS OF ANTHRANILAMIDE SCHIFF BASES: AN APPROACH TOWARD SIRTINOL

Seyed Mostafa Ebrahimi,¹ Mohammad Mahdavi,¹
Saeed Emami,² Mina Saeedi,¹ Mehdi Asadi,¹
Loghman Firoozpour,³ Mehdi Khoobi,¹ Kouros Divsalar,⁴
Abbas Shafiee,¹ and Alireza Foroumadi^{1,3}


¹Department of Medicinal Chemistry, Faculty of Pharmacy and Pharmaceutical Sciences Research Center, Tehran University of Medicinal Sciences, Tehran, Iran

²Department of Medicinal Chemistry and Pharmaceutical Sciences Research Center, Faculty of Pharmacy, Mazandaran University of Medical Sciences, Sari, Iran

³Drug Design and Development Research Center, Tehran University of Medicinal Sciences, Tehran, Iran

⁴Neuroscience Research Center, Institute of Neuropharmacology, Kerman University of Medicinal Sciences, Kerman, Iran

GRAPHICAL ABSTRACT



Abstract A novel and simple method for the green one-pot synthesis of anthranilamide Schiff bases is described. The reported Schiff bases are obtained via the reaction of isatoic anhydride, amines, and aromatic aldehydes in water at room temperature, without using any catalysts. No cyclization toward 2,3-dihydro-4(1H)-quinazolinones occurred in this method and anthranilamide Schiff bases were produced exclusively. This approach offers a green method to prepare the medicinally important Schiff base sirtinol and other bioactive anthranilamide Schiff bases.

Received July 20, 2013.

Address correspondence to Alireza Foroumadi, Department of Medicinal Chemistry, Faculty of Pharmacy and Pharmaceutical Sciences Research Center, Tehran University of Medicinal Sciences, Tehran, Iran. E-mail: aforoumadi@yahoo.com

[Supplementary materials are available for this article. Go to the publisher's online edition of Synthetic Communications® for the following free supplemental resource(s): Full experimental and spectral details.]


Keywords 2-Aryl-*N*-substituted-benzamides; isoatoic anhydride; multicomponent reactions (MCRs); Schiff base

INTRODUCTION


Isatoic anhydrides are generally used as useful building blocks in the synthesis of heterocyclic compounds such as quinazolinones, quinazolones, quinazolinediones, pyrroloquinazolones, benzimidazolones, and phthalimides with therapeutic interest.^[1] Moreover, isatoic anhydrides are also used as intermediates in the synthesis of anthranilic acid and anthranilamide derivatives with various biological activities including anticancer,^[2] VEGFR tyrosine kinase inhibitory,^[3] factor Xa inhibitory,^[4] and insecticidal activities.^[5]

Multicomponent reactions (MCRs), because of their atom economy, simple procedures, straightforward reaction design, facile execution, and convergence, are some of the best tools in synthesis of organic compounds.^[6] The utmost attribute of MCRs is the inherent formation of several bonds to construct complex molecules in one operation, ideally without isolation of intermediates.^[7] These highly step-economical reactions are particularly appealing in the context of target-oriented synthesis. They also bear the promise of novelty in terms of process and compound-related intellectual property.^[6]

In this context, three-component condensation of an isatoic anhydride, a primary amine, and an aromatic aldehyde has been widely investigated under a variety of catalysts such as *p*-TsOH,^[8] silica sulfuric acid,^[9] Al(H₂PO₄)₃,^[10] KAl(SO₄)₂ · 12H₂O (alum),^[11] [bmim]BF₄,^[12] zinc perfluorooctanoate,^[13] and gallium triflate.^[14] Using these catalysts or other conditions^[15] led to the formation of 4(3*H*)-quinazolinones **4** or 2,3-dihydro-4(1*H*)-quinazolinones **5** because of their stability (Scheme 1). To the best of our knowledge, there is no report for the production of another compound in this reaction. Following our research program on MCRs using isatoic anhydride as starting material,^[16] we found that when the latter reaction is carried out at room temperature in water, surprisingly 4(3*H*)-quinazolinones **4** or 2,3-dihydro-4(1*H*)-quinazolinones **5** are not obtained, and instead the anthranilamide Schiff bases **6** are produced (Scheme 2).



Scheme 1. Three-component reaction of isatoic anhydride **1**, amines **2**, and aldehydes **3**.



Scheme 2. One-pot synthesis of anthranilamide Schiff bases **6**.

Accordingly, herein we present a novel synthetic method for the construction of anthranilamide Schiff bases **6** via a green three-component reaction involving commercially available starting materials.

RESULTS AND DISCUSSION

It is important to note that there is only one report for the synthesis of **6**, which is prepared by a multistep reaction from anthranilic acid using extensive and dangerous reagents such as thionyl chloride and pyridine.^[17]

As shown in Scheme 2, a mixture of isatoic anhydride **1**, amine **2**, and aldehyde **3** undergoes a 1:1:1 addition reaction at ambient temperature in water to produce the corresponding anthranilamide Schiff bases **6a–m** in 65–92% yields (Table 1). All the reactions reached completion within 2–3 h. As shown in Table 1, the generality of the reaction toward various amines and benzaldehydes was explored. The results indicated that the direct three-component reactions worked well with aniline, benzylamine, furfurylamine, and substituted benzylamine. Also the reactions with a range of benzaldehydes carrying either electron-donating or withdrawing groups on the benzene ring afforded desired products in good yields.

The structures of the compounds **6** were deduced on the basis of infrared (IR), ¹H and ¹³C NMR spectroscopy, mass spectrometry, and elemental analysis. Representatively, the IR spectrum of **6h** showed absorptions at 3337 (NH), 1713 (strong, C=O) and 1633 (strong, C=N) cm⁻¹. The mass spectrum of **6h** displayed the molecular ion (M⁺) peak at *m/z* = 378. The ¹H NMR spectrum of **6h** exhibited two singlet signals recognized as arising from CH of the imine and OMe groups (δ = 8.53 and 3.56 ppm, respectively). One doublet signal at 4.50 ppm and one triplet signal at 9.34 ppm (*J* = 5.5 Hz) are due to the CH₂ and NH groups, respectively. Characteristic signals with appropriate chemical shifts and coupling constants for the 12 protons of the aromatic moieties were observed in the aromatic region of the spectrum. The ¹H decoupled ¹³C NMR spectrum of **6h** showed 18 distinct resonances, in agreement with the proposed structure.

Among various Schiff bases reported here, sirtinol (Fig. 1) attracted our attention. Sirtinol, 2-((2-hydroxynaphthalen-1-yl)methyleneamino)-*N*-(1-phenylethyl)benzamide (**6m**), acts as a specific and a direct inhibitor of sirtuin class of histone deacetylase (HDAC) activity.^[18] Synthesis of Schiff bases similar to sirtinol rarely has been discussed in the literature.^[17,18] As shown in Table 1, when the reaction of an equimolar mixture of isatoic anhydride, 1-phenylethanamine, and 2-hydroxynaphthalene-1-carbaldehyde was conducted in water at room temperature, after 3 h the expected product, sirtinol (**6m**), was obtained in good yield (87%).

Table 1. Synthesis of anthranilamide Schiff bases **6a–h** by one-pot reaction of isoic anhydride, an amine, and an aldehyde in water under catalyst-free conditions at room temperature

Entry	Amine 2	Aldehyde 3	Product 6	Mp (°C)	Yield (%)
1				120–122	84
2				124–125	85
3				144–145	86
4				142–143	65
5				179–180	92
6				165–166	83

(Continued)

Table 1. Continued

Entry	Amine 2	Aldehyde 3	Product 6	Mp (°C)	Yield (%)
7				113–114	83
8				156–157	70
9				112–113	80
10				131–132	75
11				144–146	75
12				167–168	82
13				220–222	87

^aIsolated yields.





Figure 1. Structure of a medicinally important anthranilamide Schiff base, sirtinol.



Scheme 3. Proposed mechanism for the formation of 6.

A mechanistic rationalization for this reaction is provided in Scheme 3. It is conceivable that initially isatoic anhydride **1** can convert to compound **7** via ring opening by nucleophilic attack of amine **2** to the carbonyl group, which then undergoes a decarboxylation. This generated anthranilamide could convert to the titled product **6** by simple condensation with aldehyde **3**.

A literature survey indicates that condensation of the aromatic aldehyde with the amino group of type **7** compounds in the presence of *p*-TsOH/DDQ,^[19] I₂,^[20] FeCl₃,^[21] CuCl₂,^[22] TiCl₄/Zn,^[23] chiral phosphoric acid,^[24] and ionic liquid/water^[25] undergoes cyclization to afford the dihydroquinazoline product **5**. In our method, no cyclization toward 2,3-dihydro-4(1*H*)-quinazolinones occurred and anthranilamide Schiff bases were produced exclusively.

CONCLUSION

In conclusion, we have described a simple, highly efficient, and straightforward protocol for the exclusive synthesis of anthranilamide Schiff bases **6** in water at ambient temperature without using any catalysts. Furthermore, the procedure offers several advantages including simple experimental procedure, excellent yields of the products, clean reactions, simple workup, and low cost, which make it a useful and attractive strategy in the synthesis of the medicinally important Schiff base sirtinol.

EXPERIMENTAL

Melting points were taken on a Kofler hot-stage apparatus and are uncorrected. ^1H and ^{13}C NMR spectra were recorded on Bruker FT-500, using tetramethylsilane (TMS) as an internal standard. The infrared (IR) spectra were obtained on a Nicolet Magna FT-IR 550 spectrophotometer (KBr disks). Mass spectra were recorded on an Agilent Technology (HP) mass spectrometer operating at an ionization potential of 70 eV. Elemental analyses were carried out with a Perkin-Elmer model 240-C apparatus.

Synthesis of Anthranilamide Schiff Bases 6a–m

A mixture of isatoic anhydride (1 mmol), amine (1 mmol), and aromatic aldehyde (1 mmol) in H_2O (5 mL) was stirred at room temperature for 2–3 h. After the completion of reaction (checked by thin-layer chromatography, TLC), the precipitate was filtered off and recrystallized from ethanol to give the pure compound **6**.

2-(4-Chlorobenzylideneamino)-*N*-benzylbenzamide (6a)

Yield: 84%; mp 120–122 °C; IR (KBr): 3251, 3055, 2950, 2862, 1646, 1620, 1566 cm^{-1} ; ^1H NMR (DMSO- d_6 , 500 MHz): $\delta_{\text{H}}=9.09$ (t, $J=5.6$ Hz, 1H, NH), 8.61 (s, 1H, NCH), 7.86 (dd, $J=7.7$, 1.5 Hz, 1H, ArH), 7.78 (d, $J=8.5$ Hz, 2H, ArH), 7.55 (dd, $J=7.7$, 1.5 Hz, 1H, ArH), 7.52 (d, $J=8.5$ Hz, 2H, ArH), 7.38 (dt, $J=7.7$, 1.5 Hz, 1H, ArH), 7.35–7.25 (m, 6H, ArH), 4.50 (d, $J=5.6$ Hz, 2H, NCH₂); ^{13}C NMR (DMSO- d_6 , 125 MHz): $\delta_{\text{C}}=165.9$, 161.2, 148.7, 139.0, 136.6, 134.3, 131.6, 130.6, 129.4, 129.0, 128.9, 128.3, 127.5, 126.9, 126.2, 119.2, 42.9; MS m/z (%) = 350 ($[\text{M}]^+ + 2$), 10, 348 ($[\text{M}]^+$), 30, 257 (100), 237 (61), 214 (42), 180 (41), 152 (31), 106 (80), 77 (35), 51 (14). Anal. calcd. for $\text{C}_{21}\text{H}_{17}\text{ClN}_2\text{O}$: C, 72.31; H, 4.91; N, 8.03. Found: C, 72.05; H, 4.72; N, 8.28.

SUPPORTING INFORMATION

Full experimental details and ^1H and ^{13}C NMR spectra are available via the Supplementary Content section of this article's Web page.

ACKNOWLEDGMENTS

This research was supported by grants from the research council of Tehran University of Medical Sciences and the Iran National Science Foundation (INSF).

REFERENCES

1. (a) Coppola, G. M.; Marsden, C. M. The chemistry of 2*H*-3,1-benzoxazine-2,4(1*H*)-dione (isatoic anhydride), 23 (1): Synthesis of 2-(2-propenyl)- and 2-isopropylanilines. *Heterocycl. Commun.* **1996**, 2, 301–304; (b) Kappe, T.; Stadlbauer, W. Isatoic anhydrides and their uses in heterocyclic synthesis. *Adv. Heterocycl. Chem.* **1981**, 28, 127–182.

2. (a) Congiu, C.; Cocco, M. T.; Lilliu, V.; Onnis, V. New potential anticancer agents based on the anthranilic acid scaffold: Synthesis and evaluation of biological activity. *J. Med. Chem.* **2005**, *48*, 8245–8252; (b) Cocco, M. T.; Congiu, C.; Lilliu, V.; Onnis, V. Synthesis of new *N*-(2-(trifluoromethyl)pyridin-4-yl)anthranilic acid derivatives and their evaluation as anticancer agents. *Bioorg. Med. Chem. Lett.* **2004**, *14*, 5787–5791.
3. Nakamura, H.; Sasaki, Y.; Uno, M.; Yoshikawa, T.; Asano, T.; Ban, H. S.; Fukazawa, H.; Shibuya, M.; Uehara, Y. Synthesis and biological evaluation of benzamides and benzamidines as selective inhibitors of VEGFR tyrosine kinases. *Bioorg. Med. Chem. Lett.* **2006**, *16*, 5127–5131.
4. Zhang, P.; Bao, L.; Fan, J.; Jia, Z. J.; Sinha, U.; Wong, P. W.; Park, G.; Hutchaleelaha, A.; Scarborough, R. M.; Zhu, B.-Y. Anthranilamide-based *N,N*-dialkylbenzamidines as potent and orally bioavailable factor Xa inhibitors: P4 SAR. *Bioorg. Med. Chem. Lett.* **2009**, *19*, 2186–2189.
5. Clark, D. A.; Lahm, G. P.; Smith, B. K.; Barry, J. D.; Clagg, D. G. Synthesis of insecticidal fluorinated anthranilic diamides. *Bioorg. Med. Chem.* **2008**, *16*, 3163–3170.
6. (a) Zhu, J.; Bienaymé, H. *Multicomponent Reactions*; Wiley-VCH: Weinheim, 2005; (b) Basso, A.; Banfi, L.; Riva, R.; Guanti, G. A novel highly selective chiral auxiliary for the asymmetric synthesis of L- and D- α -amino acid derivatives via a multicomponent Ugi reaction. *J. Org. Chem.* **2005**, *70*, 575–579; (c) Ramón, D. J.; Yus, M. Asymmetric multicomponent reactions (AMCRs): The new frontier. *Angew. Chem., Int. Ed.* **2005**, *44*, 1602–1634; (c) Gu, Y. Multicomponent reactions in unconventional solvents: State of the art. *Green Chem.* **2012**, *14*, 2091–2128; (c) Balalaie, S.; Arabanian, A. One-pot synthesis of tetrasubstituted imidazoles catalyzed by zeolite HY and silica gel under microwave irradiation. *Green Chem.* **2000**, *2*, 274–276.
7. (a) Tietze, L. F. Domino reactions in organic synthesis. *Chem. Rev.* **1996**, *96*, 115–136; (b) Dömling, A. Recent developments in isocyanide-based multicomponent reactions in applied chemistry. *Chem. Rev.* **2006**, *106*, 17–89; (c) Prasanna, P.; Perumal, S.; Menéndez, J. C. Chemodivergent, multicomponent domino reactions in aqueous media: L-Proline-catalyzed assembly of densely functionalized 4*H*-pyrano[2,3-*c*]pyrazoles and bispyrazolyl propanoates from simple, acyclic starting materials. *Green Chem.* **2013**, *15*, 1292–1299.
8. Baghbanzadeh, M.; Salehi, P.; Dabiri, M.; Kozehgary, G. Water-accelerated synthesis of novel bis-2,3-dihydroquinazolin-4(1*H*)-one derivatives. *Synthesis* **2006**, 344–348.
9. Salehi, P.; Dabiri, M.; Zolfigol, M. A.; Baghbanzadeh, M. A novel method for the one-pot three-component synthesis of 2,3-dihydroquinazolin-4(1*H*)-ones. *Synlett* **2005**, 1155–1157.
10. Shaterian, H. R.; Oveisi, A. R.; Honarmand, M. Synthesis of 2,3-dihydroquinazolin-4(1*H*)-ones. *Synth. Commun.* **2010**, *40*, 1231–1242.
11. Dabiri, M.; Salehi, P.; Otokesh, S.; Baghbanzadeh, M.; Kozehgary, G.; Mohammadi, A. A. Efficient synthesis of mono- and disubstituted 2,3-dihydroquinazolin-4(1*H*)-ones using $\text{KAl}(\text{SO}_4)_2 \cdot 12\text{H}_2\text{O}$ as a reusable catalyst in water and ethanol. *Tetrahedron Lett.* **2005**, *46*, 6123–6126.
12. Dabiri, M.; Salehi, P.; Baghbanzadeh, M. Ionic liquid-promoted eco-friendly and efficient synthesis of 2,3-dihydroquinazolin-4(1*H*)-ones. *Monatsh. Chem.* **2007**, *138*, 1191–1194.
13. Wang, L.-M.; Hu, L.; Shao, J.-H.; Yu, J.; Zhang, L. A novel catalyst zinc(II) perfluorooctanoate $[\text{Zn}(\text{PFO})_2]$ -catalyzed three-component, one-pot reaction: Synthesis of quinazolinone derivatives in aqueous micellar media. *J. Fluorine Chem.* **2008**, *129*, 1139–1145.
14. Chen, J.; Wu, D.; He, F.; Liu, M.; Wu, H.; Ding, J.; Su, W. Gallium(III) triflate-catalyzed one-pot selective synthesis of 2,3-dihydroquinazolin-4(1*H*)-ones and quinazolin-4(3*H*)-ones. *Tetrahedron Lett.* **2008**, *49*, 3814–3818.
15. (a) Ozaki, K.; Yamada, Y.; Oine, T.; Ishizuka, T.; Iwasawa, Y. Studies on 4(1*H*)-quinazolinones, 5: Synthesis and anti-inflammatory activity of 4(1*H*)-quinazolinone derivatives. *J. Med. Chem.* **1985**, *28*, 568–576; (b) Sadanandam, Y. S.; Reddy, K. R. M.; Rao,

- A. B. Synthesis of substituted 2,3-dihydro-1-(β -phenylethyl)-2-aryl and 2,3-diaryl-4(1*H*)-quinazolinones and their pharmacological activities. *Eur. J. Org. Chem.* **1987**, *22*, 169–173; (c) Salehi, P.; Dabiri, M.; Zolfigol, M. A.; Baghbanzadeh, M. A new approach to the facile synthesis of mono- and disubstituted quinazolin-4(3*H*)-ones under solvent-free conditions. *Tetrahedron Lett.* **2005**, *46*, 7051–7053; (d) Lingaiah, B. V.; Ezekiel, G.; Yakaiah, T.; Reddy, G. V.; Rao, P. S. Nafion-H: An efficient and recyclable heterogeneous catalyst for the one-pot synthesis of 2,3-disubstituted 4-(3*H*)-quinazolinones under solvent-free microwave irradiation conditions. *Synlett* **2006**, 2507–2509.
16. (a) Mahdavi, M.; Asadi, M.; Saeedi, M.; Rezaei, Z.; Moghbel, H.; Foroumadi, A.; Shafiee, A. Synthesis of novel 1,4-benzodiazepine-3,5-dione derivatives: Reaction of 2-aminobenzamides under Bargellini reaction conditions. *Synlett* **2012**, *23*, 2521–2525; (b) Mahdavi, M.; Asadi, M.; Saeedi, M.; Tehrani, M. H.; Mirfazli, S. S.; Shafiee, A.; Foroumadi, A. Green synthesis of new boron-containing quinazolinones: Preparation of benzo[*d*][1,3,2]diazaborinin-4(1*H*)one derivatives. *Synth. Commun.* **2013**, *43*, 2936–2942; (c) Asadi, M.; Ebrahimi, M.; Mahdavi, M.; Saeedi, M.; Ranjbar, P. R.; Yazdani, F.; Shafiee, A.; Foroumadi, A. Reaction of isatoic anhydride, amine, and *N,N'*-dialkyl carbodiimides under solvent-free conditions: New and efficient synthesis of 3-alkyl-2-(alkylamino) quinazolin-4(3*H*)-ones. *Synth. Commun.* **2013**, *43*, 2385–2392.
 17. Liang, Y.; Su, B.; Zhao, J.; Sun, W. The synthesis of new asymmetric double Schiff bases containing a new *o*-amino benzoic acid derivative. *Synth. Commun.* **2004**, *34*, 3235–3242.
 18. Mai, A.; Massa, S.; Lavu, S.; Pezzi, R.; Simeoni, S.; Ragno, R.; Mariotti, F. R.; Chiani, F.; Camilloni, G.; Sinclair, D. A. Design, synthesis, and biological evaluation of sirtinol analogues as class III histone/protein deacetylase (sirtuin) inhibitors. *J. Med. Chem.* **2005**, *48*, 7789–7795.
 19. Shaabani, A.; Maleki, A.; Mofakham, H. Click reaction: Highly efficient synthesis of 2,3-dihydroquinazolin-4(1*H*)-ones. *Synth. Commun.* **2008**, *38*, 3751–3759.
 20. Bhat, B. A.; Sahu, D. P. One-pot synthesis of 4(3*H*)-quinazolinones. *Synth. Commun.* **2004**, *34*, 2169–2176.
 21. Wang, G.-W.; Miao, C.-B.; Kang, H. Benign and efficient synthesis of 2-substituted 4(3*H*)-quinazolinones mediated by iron(III) chloride hexahydrate in refluxing water. *Bull. Chem. Soc. Jpn.* **2006**, *79*, 1426–1430.
 22. Abdel-Jalil, R. J.; Voelter, W.; Saeed, M. A novel method for the synthesis of 4(3*H*)-quinazolinones. *Tetrahedron Lett.* **2004**, *45*, 3475–3476.
 23. Shi, D.; Rong, L.; Wang, J.; Zhuang, Q.; Wang, X.; Hu, H. Synthesis of quinazolin-4(3*H*)-ones and 1,2-dihydroquinazolin-4(3*H*)-ones with the aid of a low-valent titanium reagent. *Tetrahedron Lett.* **2003**, *44*, 3199–3201.
 24. (a) Cheng, X.; Vellalath, S.; Goddard, R.; List, B. Direct catalytic asymmetric synthesis of cyclic amins from aldehydes. *J. Am. Chem. Soc.* **2008**, *130*, 15786–15787; (b) Rueping, M.; Antonchick, A. P.; Sugiono, E.; Grenader, K. Asymmetric Brønsted acid catalysis: Catalytic enantioselective synthesis of highly biologically active dihydroquinazolinones. *Angew. Chem., Int. Ed.* **2009**, *48*, 908–910.
 25. Chen, J.; Su, W.; Wu, H.; Liu, M.; Jin, C. Eco-friendly synthesis of 2,3-dihydroquinazolin-4(1*H*)-ones in ionic liquids or ionic liquid–water without additional catalyst. *Green Chem.* **2007**, *9*, 972–975.