

Baseline insecticide susceptibility data of *Phlebotomus papatasi* in Iran

Z. Saeidi¹, H. Vatandoost¹, A.A. Akhavan¹, M.R. Yaghoobi-Ershadi¹, Y. Rassi¹,
M.H. Arandian² & R. Jafari²

¹Department of Medical Entomology and Vector Control, School of Public Health, Tehran University of Medical Sciences, Tehran; ²Esfahan Health Research Station, National Institute of Health Research, Tehran University of Medical Sciences, Esfahan, Iran

ABSTRACT

Background & objectives: Phlebotomine sandflies (Diptera: Psychodidae) play main role in the transmission of different forms of leishmaniasis in the world. *Phlebotomus papatasi* is the main vector of zoonotic cutaneous leishmaniasis (ZCL) in Iran. There are several control measures for vector control using different classes of insecticides. The aim of this study was to breed the sandflies which were collected from a hyperendemic focus of the disease in central Iran in the laboratory condition and to determine its baseline susceptibility to commonly used insecticides.

Methods: Sandflies were collected from the field and were reared in insectary. Susceptibility tests were carried out on their generation. Baseline susceptibility of sandflies to DDT and pyrethroids was evaluated based on LT₅₀ values. A total of 1305 specimens were tested using different time intervals. The LT₅₀ and LT₉₀ values were measured according to the WHO standard tests. The results were plotted using probit analysis and regression lines.

Results: The results against female sandflies revealed the LT₅₀ values of 1312.66, 253.66, 36.04, 9.38 and 6 sec to DDT (4%), permethrin (0.75%), deltamethrin (0.1%), cyfluthrin (0.15%) and lambda-cyhalothrin (0.05%), respectively. The figures for male sandflies were 1200.97, 310.10, 18.63, 6.08 and 0.77 sec respectively to the above insecticides.

Conclusion: The results of this study could help to provide a clue for implementation of currently used insecticides. Furthermore, a specific guideline is needed for monitoring and evaluation of insecticide susceptibility test against sandflies.

Key words Control; insecticides; Iran; *Phlebotomus papatasi*; zoonotic cutaneous leishmaniasis

INTRODUCTION

Leishmaniases are a group of vector-borne diseases transmitted by phlebotomine sandfly and caused by protozoan parasite of genus *Leishmania*^{1–3}. Leishmaniasis is endemic in 98 countries or territories⁴ — most of them are developing countries in tropical and subtropical parts of the world. In these areas, about 350 million population is at risk with about 12 million reported infections and about 60,000 deaths annually³. It is estimated that 1.5–2 million people are infected per year with 500,000 new cases for visceral leishmaniasis (VL) and 1–1.5 million for cutaneous leishmaniasis (CL), and of these only 600,000 cases are officially declared. The disease burden is estimated at 2,356,000 (946,000 in men and 1,410,000 in women) disability adjusted life years (DALYs); that show a significant ranking among communicable diseases⁵. Ninety percent of CL cases occur in 7 countries (Afghanistan, Algeria, Brazil, Iran, Peru, Saudi Arabia and Syria) and 90% of VL cases occur in rural and suburban areas in 5 countries (Bangladesh, India, Nepal, Sudan and Brazil)². Different laboratories tried to rear sandflies in different parts of the

world, each of them used special methods for feeding⁶, temperature and other conditions^{7–9}. Some of them encountered with some difficulties in rearing process like fungal contamination¹⁰.

There are 56 species (32 *Phlebotomus* and 24 *Sergentomyia*) of phlebotomine sandflies in Iran but *Phlebotomus papatasi* is the main vector of ZCL¹¹. Some researchers carried out susceptibility of sandflies in different parts of the world. They reported *P. papatasi* resistance to DDT in northern Bihar (India)¹² and Turkey^{1,11}. In some parts of Rajasthan, *P. papatasi* was resistant to DDT but susceptible to dieldrin, malathion, fenitrothion and propoxur¹³. In other parts of Rajasthan this species was found resistant to DDT, dieldrin and propoxur while it was susceptible to malathion, fenitrothion and permethrin¹³ while another study in Bihar showed that *P. argentipes* was resistant to DDT (4%) but susceptible to deltamethrin (0.05%)¹⁴. Italian populations of *P. perniciosus* and *P. papatasi* from Campania region and from Rome, respectively, were susceptible to the insecticides — (DDT (2%), lambda-cyhalothrin (0.06%) and permethrin (0.2%) as compared with the reference strain used¹⁵. In countries

where there are vector control operations for malaria, sandflies population could also be controlled. However, after stopping the use of insecticide, the number of leishmaniasis cases was increased^{1,16}. In different areas of Iran, researchers carried out several tests. Results showed that in the ACL foci in northeastern part, *P. sergenti* was susceptible to deltamethrin¹⁷ and DDT (4%) in Esfahan¹⁸, and *P. papatasi* was susceptible to deltamethrin in Borkhar county, Esfahan province¹⁹, susceptible to DDT in Badrood county²⁰, susceptible to DDT (4%), permethrin (0.25%), propoxur (0.1%) in Sabzevar²¹, susceptible to DDT (4%) in Orzouiyeh county, Kerman province^{22,11} and susceptible to DDT (4%) in Jarghuyr county, Esfahan province and tolerant to DDT (4%) but susceptible to permethrin (0.75%), deltamethrin (0.1%), cyfluthrin (0.15%), and lambda-cyhalothrin (0.05%)¹¹. Due to lack of enough information about susceptibility status of *P. papatasi* to recommended insecticides by WHO after testing the wild strain, we decided to test the laboratory-reared one against DDT and some current used pyrethroids²³.

MATERIAL & METHODS

Study area

This survey was conducted during the summer 2010 and sandflies were originated from the rural district of Badrood, Natanz county, Esfahan province, central Iran. This area is located in the foothills of the Karkas Mountains (altitude 3895 m). The area has a semi-desert climate. All the sandflies were collected from Matin Abad tourist camp that was situated in altitude of 978 m (33°45' N, 51°59' E).

Sandfly collection

Adults of *P. papatasi* were collected using hand aspirator and torch. The adults were caught from outdoor during dawn and dusk from parked car near to their breeding places. Collected sandflies were transferred to a cage with a hanging piece of wet cloth for supplying suitable humidity and were fed by a small amount of sucrose solution soaked cotton. All of them were placed in a plastic bag to remain wet and to maintain stable temperature situation. Subsequently, the cages were transported to Tehran sandfly insectary in the School of Public Health, Tehran University of Medical Sciences for sandfly maintenance and rearing²⁴. Temperature was maintained by automated electric heaters and photoperiod of 14/10 H D/L was maintained in the insectary.

Sandfly rearing

After resting, fed and gravid wild-caught female

adults were separated by aspirator and were released into individual pots according to Killick-Kendrick and Killick-Kendrick method²⁵ and were fed with honey solution (50%) and saturated sucrose. The pots were checked daily for hatching the eggs. The larvae (L₁) were fed with larval food, complex of rabbit food (palette) and rabbit feces without liver powder. In pupal stage the exuviate of L₄ and the tails exist at the end of pupae body. Before adding food for larvae, small amount of autoclaved sea sand was added. This is to prevent the larval movement in case of fungal infestation and were checked daily to take out fungi. Emerged adults were released in a new cage with wet cloth and sucrose solution (20%). Then the 3–10 days old adults were tested in a standard WHO susceptibility test method as described for mosquitoes. Identification of the adult females and males, was done based on Modi and Tesh method²⁶. There are lot of problems in sandfly rearing such as fungal contamination, mite contamination, cannibalism and some anomaly in sandfly bodies. For instance, fungal contamination can trap larvae, especially I instar. For precaution of fungal growth, all materials and instruments were autoclaved at 121°C for 20 min. All the larval pots were disinfected. For mite protection, the insectary must has a routine washing program with warm water. Sometimes cannibalism could be observed in some pots especially at the I instar larvae. Sometime abnormality can be found in bodies at different stages.

Susceptibility tests

After some resting and feeding by sucrose solution (20%), sandflies were tested according to the standard method of WHO²⁷. During the tests, the sandflies were released into the exposure tubes at different time intervals and then the mortality was counted after 24 h recovery period. During the holding time, the insects were supplied with cotton pad soaked in sucrose solution (20%). All the mortalities were corrected according to the results of control with Abbott's correction²⁸. All the tests were excluded when the mortality was > 20% in the control group. After each test, all the dead and alive sandflies were transferred to 75% alcohol separately for mounting in Pouri's medium for species identification. Males and females were counted separately. Females and males at least 24 h after mounting, were identified using valid key^{29, 30} and if there were any other species, all of them were excluded from the tests. Females of *P. papatasi* were identified by regular segments in spermatheca and network in pharyngeal armature.

The exposure time interval was between 7 and 3600 sec. At least 5 interval times were used to gain the mor-

tality between 5 and 95%. In each exposure time at least 4 replicates were used comprising 50–100 sandflies depending on the availability and the same age of the adults. The susceptibility tests were carried out on 1305 laboratory-reared *P. papatasi* (691 females and 614 males)

Procurement of insecticide papers and their concentration

Impregnated papers (DDT 4%, permethrin 0.75%, deltamethrin 0.1%, cyfluthrin 0.15%, lambda-cyhalothrin 0.05%) were procured from collaborating center of WHO in Malaysia.

Data analysis

The exposure time versus probit mortality were used according to Finney 1971. The Excel was used for data entering. HG4 software was used for drawing the graphs.

RESULTS

The results of tests against the laboratory-reared females and males are shown in Figs. 1 and 2, respectively. The results of susceptibility test against laboratory-reared




Fig. 1: Probit regression lines of different insecticides against laboratory-reared *P. papatasi* females.




Fig. 2: Probit regression lines of different insecticides against laboratory-reared *P. papatasi* males.




Fig. 3: LT_{50} values of different insecticides against laboratory-reared males and females *P. papatasi*.

female *P. papatasi* revealed LT_{50} values of 1312.66, 253.66, 36.47, 9.38, 6 sec to DDT (4%), permethrin (0.75%), deltamethrin (0.1%), cyfluthrin (0.15%) and lambda-cyhalothrin (0.05%) respectively. This data for males were 1200.97, 310.10, 18.63, 6.08 and 0.77 sec respectively to the above insecticides. The results showed that males were more susceptible than females to all the insecticides tested at LT_{50} level (Fig. 3).

DISCUSSION & CONCLUSION

The main measures for ZCL control in Iran are rodent control operation, using impregnated bednets and curtains with pyrethroids, the use of repellents, indoor residual spraying (IRS), and health education to the community during complex emergency situations, leishmanization is also recommended. Owing to the lack of information about the susceptibility of sandflies to different WHO recommended insecticides, the present study was undertaken. WHO susceptibility test recommended for mosquitoes was followed. The discriminative dose of DDT and pyrethroids were used in the tests. In accordance to WHO report³¹, the bioassay results for malaria vectors were summarized in three resistant classes: susceptible with mortality rate 98–100%; possibly resistant (tolerant) with mortality rate between 97 and 80%; and resistant with mortality rate < 80%. Results of the test against laboratory-reared for females of *P. papatasi* at the LT_{50} values exhibited that males are more susceptible than females to all of the insecticides tested. Results of our tests against susceptible *P. papatasi* at LT_{50} level revealed that the females needs more time to be killed at the same concentration than males.

The time for mortality of sandflies was more than the pyrethroids. The high LT_{50} level of the vector to DDT is attributed to the long-term use of insecticides for malaria vector control in the region (from 1953 for up to 5 yr DDT

was applied as indoor residual spraying for malaria control in Badrood) that transmitted genetically to their progeny. In Esfahan province, several herbicides, fungicides and insecticides have been used for agriculture and veterinary pest control, including diazinon, malathion, azinphosmethyl, fenitrothion, metasytox, permethrin, carbaryl and cypermethrin. There are several reports of susceptibility of *Leishmania* vectors to different insecticides in Iran. For example, *P. papatasi* found susceptible to deltamethrin in Borkhar district, Esfahan province¹⁹, and susceptible to DDT in Badrood district²⁰. In Sabzevar district susceptible to DDT (4%), permethrin (0.25%) and propoxur (0.1%)²¹; and in Orzuie district in Kerman province susceptible to DDT (4%)²². *Phlebotomus sergenti* found susceptible to deltamethrin in Mashhad, northeast of Iran¹⁷ and susceptible to DDT (4%) in Esfahan province¹⁸. There are several reports of reservoir hosts and their control in Iran^{32,33}. As we mentioned earlier¹¹, we recommend the same procedure in different parts of the world to pool the results and reach the unique conclusion about criteria for susceptibility status against *P. papatasi*. From the pooled results World Health Organization is able to provide a specific guideline for sandfly and this guideline will help the countries for monitoring and evaluation of insecticide resistance for implementation of control measures.

ACKNOWLEDGEMENTS

The authors would like to express their appreciation to Tehran University of Medical Sciences for kindly providing financial support to this research.

REFERENCES

- Lane RP. Sandflies (Phlebotominae) In: Lane RP, Crosskey RW, editors. *Medical insects and arachnids*. London: Chapman and Hall 1993; p. 78–119.
- Gramiccia M, Gradoni L. The current status of zoonotic leishmaniasis and approaches to disease control. *Int J Parasitol* 2005; 35: 1169–80.
- Leishmaniasis, the disease and its epidemiology. Available from: http://www.who.int/leishmaniasis/disease_epidemiology/en/ (2010).
- Report of a meeting of the WHO expert committee on the control of Leishmaniasis. Geneva: World Health Organization 2010.
- Report of the consultative meeting on cutaneous leishmaniasis. Geneva: World Health Organization 2008. WHO/HTM/NTD/IDM/2008.7.
- Harre JG, Dorsey KM, Armstrong KL, Burge JR, Kinnamon KE. Comparative fecundity and survival rates of *Phlebotomus papatasi* sandflies membrane fed on blood from eight mammal species. *Med Vet Entomol* 2001; 15: 189–96.
- Chelbi I, Zhioual E. Biology of *Phlebotomus papatasi* (Diptera: Psychodidae) in the laboratory. *J Med Entomol* 2007; 44(4): 597–600.
- Kasap OE, Alten B. Laboratory estimation of degree-day developmental requirements of *Phlebotomus papatasi* (Diptera: Psychodidae). *J Vector Ecol* 2005; 30 (2): 328–33.
- Kasap OE, Alten B. Comparative demography of the sandfly *Phlebotomus papatasi* (Diptera: Psychodidae) at constant temperatures. *J Vector Ecol* 2006; 31 (2): 378–85 .
- Marchais R, Bouchet F, Bouchet P. Laboratory rearing of *Phlebotomus perniciosus* (Diptera: Psychodidae) and fungal growth problems. *Parassit* 1991; 33 (Suppl): 393–7.
- Saeidi Z, Vatandoost H, Akhavan AA, Yaghoobi-Ershadi MR, Arandian MH, Rassi Y, Jafari R, Sheikh Z, Sanei Dehkordi AR. Baseline susceptibility of a wild strain of *Phlebotomus papatasi* (Diptera: Psychodidae) to DDT and pyrethroids in an endemic focus of zoonotic cutaneous leishmaniasis in Iran. *Pest Manag Sci* 2012; 68(5): 669–75.
- Joshi GC, Kaul SM, Wattal BL. Susceptibility of sandflies to organochlorine insecticides in Bihar, India: Further reports. *J Com Dis* 1979; 11: 209–13 .
- Singh KV, Bansal SK. Insecticide susceptibility of *Phlebotomus papatasi* to organochlorine, organophosphate and carbamate compounds in some arid areas of western Rajasthan. *Indian J Med Res* 1996; 103: 91–3.
- Dinesh DS, Kumar V, Das P. Efficacy of DDT and deltamethrin on sandfly population in an endemic village of kala-azar, Bihar, India. *7th International Symposium on Phlebotomine Sandflies (ISOPS7)*, April 25–30, 2011. Kusadasi, Turkey: Pine Bay Holiday Resort 2011; p. 112.
- Maroli M, Cianchi T, Bianchi R, Khoury C. Testing insecticide susceptibility of *Phlebotomus perniciosus* and *P. papatasi* (Diptera: Psychodidae) in Italy. *Ann Ist Super Sanità* 2002; 38(4): 419–23.
- Seyedi-Rashti MA, Nadim A. Attempts to control zoonotic cutaneous leishmaniasis in Isfahan area, Iran. *9th International Congress for Tropical Medicine and Malaria, Athens 1974*; p. 135.
- Moosa-Kazemi SH, Yaghoobi-Ershadi MR, Akhavan AA, Abdoli H, Zahraei-Ramazani AR, Jafari R, et al. Deltamethrin-impregnated bednets and curtains in an anthroponotic cutaneous leishmaniasis control program in northeastern Iran. *Annals Saudi Med* 2007; 27(1): 6–12.
- Zahraei-Ramazani AR, Yaghoobi-Ershadi MR, Akhavan AA, Abdoli H, Jafari R, Jalali-Zand AR, et al. Some ecological aspects of Phlebotominae sandflies (Diptera: Psychodidae) in an endemic focus of anthroponotic cutaneous leishmaniasis of Iran. *J Entomol* 2008; 5(1): 17–23.
- Yaghoobi-Ershadi MR, Moosa-Kazemi SH, Zahraei-Ramazani AR, Jalali-Zand AR, Akhavan AA, Arandain MH, et al. Evaluation of deltamethrin-impregnated bednets and curtains for control of zoonotic cutaneous leishmaniasis in a hyperendemic area of Iran. *Bull Soc Pathol Exot* 2006; 99(1): 43–8..
- Yaghoobi-Ershadi MR, Akhavan AA. Entomological survey of sandflies (Diptera: Psychodidae) in a new focus of zoonotic cutaneous leishmaniasis in Iran. *Acta Trop* 1999; 73(3): 321–6.
- Yaghoobi-Ershadi MR, Akhavan AA, Zahraei-Ramazani AR, Abai MR, Ebrahimi B, Vafaei-Nezhad R, et al. Epidemiological study in a new focus of cutaneous leishmaniasis in the Islamic Republic of Iran. *Eastern Mediterr Health J* 2003; 9(4): 816–26.
- Akhavan AA, Yaghoobi-Ershadi MR, Hasibi F, Jafari R, Abdoli H, Arandian MH, et al. Emergence of cutaneous leishmaniasis due to *Leishmania major* in a new focus of southern Iran. *Iran J Arthropod-Borne Dis* 2007; 1(1): 1–8.
- Test procedures for insecticide resistance monitoring in malaria vectors, bio-efficacy and persistence of insecticides on treated

- surfaces. Geneva: World Health Organization. WHO/CDS/CPC/MAL/98.12. Available from: http://wholibdoc.who.int/hq/1998/WHO_CDS_CPC_MAL_98.12.pdf.
24. Yaghoobi-Ershadi MR, Shirani-Bidabadi L, Hanafi-Bojd AA, Akhavan AA, Zeraati H. Colonization and biology of *Phlebotomus papatasi*, the main vector of cutaneous leishmaniasis due to *Leishmania major*. *Iran J Public Health* 2007; 36(3): 21–6.
 25. Killick-Kendrick M, Killick-Kendrick R. The initial establishment of sandfly colonies. *Parassitologia* 1991; 33 (Suppl): 315–20.
 26. Modi GB, Tesh RB. A simple technique for mass rearing *Lutzomyia longipalpis* and *Phlebotomus papatasi* (Diptera: Psychodidae) in the laboratory. *J Med Entomol* 1983; 20: 568–9.
 27. *Instructions for determining the susceptibility or resistance of adult mosquito to organochlorine, organophosphorine and carbamate insecticides-diagnostic test*. WHO/VBC/81.806, 1981.1.
 28. Abbott WS. A method of computing the effectiveness of an insecticide. *J Econ Entomol* 1925; 18: 265–7.
 29. Theodor O, Mesghali A. On the Phlebotomine of Iran. *J Med Entomol* 1964; 1: 285–300.
 30. Seyedi-Rashti MA, Nadim A. The genus *Phlebotomus* (Diptera: Psychodidae: Phlebotominae) of the countries of the Eastern Mediterranean region. *Iran J Public Health* 1992; 21 (1–4): 11–50.
 31. *Report of the WHO informal consultation. Test procedures for insecticide resistance monitoring in malaria vectors, bioefficacy and persistence of insecticides on treated surfaces*. WHO/CDS/CPC/MAL/98.12. Geneva: World Health Organization 1998.
 32. Rassi Y, Javadian E, Amin M, Rafizadeh S, Vatandoost H, Motazedian H. *Meriones libycus* is the main reservoir of zoonotic cutaneous leishmaniasis in south Islamic Republic of Iran. *Eastern Mediterr Health J* 2006; 12(3–4): 474–7.
 33. Rassi Y, Gassemi MM, Javadian E, Rafizadeh S, Motazedian H, Vatandoost H. Vectors and reservoirs of cutaneous leishmaniasis in Marvdasht district, southern Islamic Republic of Iran. *Eastern Mediterr Health J* 2007; 13(3): 686–93.

Correspondence to: Prof. Hassan Vatandoost and Dr Amir Ahmad Akhavan, Department of Medical Entomology and Vector Control, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran.

E-mail: hvatandoost1@yahoo.com, vatando@tums.ac.ir; aaakhavan@yahoo.com

Received: 1 September 2012

Accepted in revised form: 31 January 2013